# Index to Volume 69 Author Index

Abe, J. See A. Suzuki, 309

Adalsteins, A. E. See W. E. Werner, 535

Addo, K., and Y. Pomeranz. Effects of lipids and emulsifiers on alveograph characteristics, 6

Aguilar, F. See J.-L. Jane, 96

Aibara, S., K. Hisaki, and K. Watanabe. COMMUNICATION TO THE EDITOR: Effects of high-voltage electric field treatment on wheat dough and breadmaking properties, 465

Akesson, G. See B. Ekstrand, 379

Aktan, B., and K. Khan. Effect of high-temperature drying of pasta on quality parameters and on solubility, gel electrophoresis, and reversedphase high-performance liquid chromatography of protein components, 288

Allen, H. M. See J. R. Oliver, 546

Al-Ruqaie, I., and K. Lorenz. Alkylresorcinols in extruded cereal brans,

Andrews, S. B., T. J. Siebenmorgen, and A. Mauromoustakos. Evalua-tion of the McGill No. 2 rice miller, 35

Asami, K. See M. Ito, 325

Autio, K., and E. Sinda. Frozen doughs: Rheological changes and yeast viability, 409

\_\_\_\_. See Y. Mälkki, 647 Autran, J.-C. See P. Scheromm, 664

Bailey, T. B. See S. R. Fox, 191 Bakker-Arkema, F. W. See R. G. Moreira, 390

Batey, I. L. See R. B. Gupta, 125 Bekes, F., and P. W. Gras. COMMUNICATION TO THE EDITOR: Demonstration of the 2-gram mixograph as a research tool, 229 BeMiller, J. N. See J. E. Fannon, 284, 456

Bergoin, A. See P. Scheromm, 664
Bern, C. J. See M. O. Racker, 66
Bertrand, D., C. Le Guernevé, D. Marion, M. F. Devaux, and P. Robert. Description of the textural appearance of bread crumb by video image analysis, 257

Betschart, A. A. See T. S. Kahlon, 485

. See B. E. Knuckles, 198

Bhatty, R. S. B-Glucan content and viscosities of barleys and their roller-

milled flour and bran products, 469
Biemer, P. See F. R. Del Valle, 156
Biliaderis, C. G., M. S. Izydorczyk, O. M. Lukow, and W. Bushuk. NOTE: Pentosans in flours of 1B/1R translocation wheats, 226

Blakeney, A. B. See J. R. Oliver, 546 Bock, M. A. See F. R. Del Valle, 156 Bourges, H. See F. R. Del Valle, 156 Bouzaher, A. See A. J. Peplinski, 397

Branlard, G., D. Khelifi, and G. Lookhart. NOTE: Identification of some wheat proteins separated by a two-step acid polyacrylamide gel electrophoresis and sodium dodecyl sulfate-polyacrylamide gel electrophoresis technique, 677

Bruinsma, B. L. See K. F. Finney, 347 Brümmer, J.-M. See B. Fretzdorff, 266 Bushuk, W. See C. G. Biliaderis, 226

See L. Gao, 452 See Y. Inoue, 423 See A. Kawka, 92

. See M. Kazemie, 103, 105

Callaghan, P. T. See S. L. Umbach, 637

Camire, M. E. See S. I. Flint, 444 Carr, M. E., S. Kim, K. J. Yoon, and K. D. Stanley. Graft polymerization of cationic methacrylate, acrylamide, and acrylonitrile monomers onto starch by reactive extrusion, 70

Case, S. É., D. D. Hamann, and S. J. Schwartz. Effect of starch gelat-inization on physical properties of extruded wheat- and corn-based products, 401

Castro, M. See B. R. Hamaker, 461

Chambers, E., IV. See C.-Y. Chang, 556 Champagne, E. T., and R. J. Kron, Sr. Stabilizing brown rice to lipolytic hydrolysis by ethanol vapors, 152

Chandrashekar, A. See R. Kavitha, 440

See D. L. Rom, 178

Chang, C.-Y., and E. Chambers IV. Flavor characteristics of breads made from hard red winter wheat and hard white winter wheat, 556 Chauhan, G. S., N. A. M. Eskin, and R. Tkachuk. Nutrients and anti-

nutrients in quinoa seed, 85

Chedid, L. L., and J. L. Kokini. Influence of protein addition on rheological properties of amylose- and amylopectin-based starches in excess Chen, J., K. Khan, D. R. Shelton, and B. L. D'Appolonia. Isolation and fractionation of carbohydrate-containing proteins from wheat gluten, 475

., D. R. Shelton, B. L. D'Appolonia, and K. Khan. Structural characterization of the carbohydrate portion of a glycopeptide from wheat

gluten, 481 .. See J.-L. Jane, 528

Chen, J.-F. See J.-L. Jane, 60

Chinachoti, P. See A. Nussinovitch, 678

See P. A. Rao, 613 Chiu, M. M. See T. S. Kahlon, 485

See B. E. Knuckles, 198 Chow, F. I. See T. S. Kahlon, 485

Chung, O. K. See A. Xu, 495, 502 Clark, D. See B. R. Hamaker, 461

Clarke, J. M. See T. N. McCaig, 671

Clements, R. L. Effects of metal-complexing agents on water binding by gluten, 315

Cox, T. S. See H. Dong, 132

Curran, S. P. See K. Shelke, 141, 145

Czuchajowska, Z., B. Paszczynska, and Y. Pomeranz. Functional proper-

ties of psyllium in wheat-based products, 516
\_\_\_\_, J. Szczodrak, and Y. Pomeranz. Characterization and estimation of barley polysaccharides by near-infrared spectroscopy. I. Barleys, starches, and B-D-glucans, 413

See J. Szczodrak, 419

D'Appolonia, B. L. See J. Chen, 475, 481
\_\_\_\_. See D. P. Puhr, 582

Davis, E. A. See L. L. P. Lambert, 303
See S. L. Umbach, 637

Delost-Lewis, K., K. Lorenz, and R. Tribelhorn. Puffing quality of experi-

mental varieties of proso millets (*Panicum miliaceum*), 359

Del Valle, F. R., M. Escobedo, A. Sanchez-Marroquin, H. Bourges, M. A. Bock, and P. Biemer. Nitrogen balance in infants fed formulas

containing amaranth or a soy-oats formula, 156 Delwiche, S. R., R. E. Pitt, and K. H. Norris. NOTE: Sensitivity of near-infrared absorption to moisture content versus water activity in

starch and cellulose, 107
DePauw, R. M. See T. N. McCaig, 671
De Stefanis, V. A. NOTE: Analysis of potassium bromate in flour, dough, and bread, 683

Devaux, M. F. See D. Bertrand, 257

Dexter, J. E. See R. R. Matsuo, 27

—. See S. J. Symons, 137

Dintzis, F. R., J. Lehrfeld, T. C. Nelsen, and P. L. Finney. Phytate content of soft wheat brans as related to kernel size, cultivar, location, and milling and flour quality parameters, 577

Donelson, J. R. See G. S. Gaines, 120
Dong, H., R. G. Sears, T. S. Cox, R. C. Hoseney, G. L. Lookhart, and M. D. Shogren. Relationships between protein composition and mixograph and loaf characteristics in wheat, 132

Dorsey-Redding, C. See S. R. Fox, 191
Dronzek, B. L. See N. D. G. White, 182
Duarte, A. M. A. See F. C. Strong III, 659

Eckhoff, S. R. See A. H. Mistry, 82, 202, 296, 433

Eckhoft, S. R. See A. H. Mistry, 82, 202, 296, 433

—. See R. Ruan, 600

Edwards, N. M. See R. R. Matsuo, 27

Ekstrand, B., I. Gangby, and G. Åkesson. Lipase activity in oats—distribution, pH dependence, and heat inactivation, 379

Eliasson, A.-C. See L. Lindahl, 30, 542

—. See K. Mani, 222

Escobedo, M. See F. R. Del Valle, 156

Eskin, N. A. M. See G. S. Chauhan, 85

Fannon, J. E., and J. N. BeMiller. Structure of corn starch paste and granule remnants revealed by low-temperature scanning electron microscopy after cryopreparation, 456 , R. J. Hauber, and J. N. BeMiller. Surface pores of starch granules,

Faubion, J. M. See K. Shelke, 141, 145 Finney, K. F., B. L. Bruinsma, and O. Natsuaki. NOTE: Copper (II) vs. zinc inorganic salts as oxidizers in breadmaking, 347 Finney, P. L. See F. R. Dintzis, 577

See C. S. Gaines, 115, 120

Flint, S. I., and M. E. Camire. Recovery of lignin during nonstarch polysaccharide analysis, 444

Fox, S. R., L. A. Johnson, C. R. Hurburgh, Jr., C. Dorsey-Redding,

and T. B. Bailey. Relations of grain proximate composition and physical

properties to wet-milling characteristics of maize, 191
Fretzdorff, B., and J.-M. Brümmer. Reduction of phytic acid during

breadmaking of whole-meal breads, 266
Frey, K. J. See H. S. Lim, 262
Frye, A. M., and C. S. Setser. Optimizing texture of reduced-calorie yellow layer cakes, 338

Gahlawat, P., and S. Sehgal. NOTE: Phytic acid, saponins, and polyphenols in weaning foods prepared from oven-heated green gram and

Gaines, C. S., A. Kassuba, and P. L. Finney. Instrumental measurement of cookie hardness. I. Assessment of methods, 115

and J. R. Donelson. Instrumental measurement of cookie hardness. II. Application to product quality variables, 120

See F. R. Huebner, 148 Gangby, I. See B. Ekstrand, 379

Gao, L., P. K. W. Ng, and W. Bushuk. Structure of glutenin based on farinograph and electrophoretic results, 452

Garcia, H. H. See B. R. Hamaker, 461 Gelroth, J. A. See G. S. Ranhotra, 75, 623 Gilman, R. See B. R. Hamaker, 461 Glaser, B. K. See G. S. Ranhotra, 75, 623

Glatz, B. A. See M. O. Raeker, 66 Gomez, M. H., J. K. Lee, C. M. McDonough, R. D. Waniska, and L. W. Rooney. Corn starch changes during tortilla and tortilla chip

processing, 275 Gonzales, A. E. See B. R. Hamaker, 461 Gordon, J. See L. L. P. Lambert, 303

See S. L. Umbach, 637
Gras, P. W., and L. O'Brien. Application of a 2-gram mixograph to early generation selection for dough strength, 254 See F. Bekes, 229

Greenblatt, G. A. See C. F. Morris, 467 Greene, L. W. See S. O. Serna-Saldivar, 78 Grove, M. J. See C. A. Knutson, 436 Gupta, P. P. See N. Singh, 55

Gupta, R. B., I. L. Batey, and F. MacRitchie. Relationships between protein composition and functional properties of wheat flours, 125 Guritno, N. See R. K. Newman, 240

 Haarasilta, S. See A. Törrönen, 355
 Hamaker, B. R., C. Valles, R. Gilman, R. M. Hardmeier, D. Clark,
 H. H. Garcia, A. E. Gonzales, I. Kohlstad, M. Castro, R. Valdivia, Rodriguez, and M. Lescano. NOTE: Amino acid and fatty acid profiles of the Inca peanut (Plukenetia volubilis), 461

Hamann, D. D. See S. E. Case, 401

Hamer, R. J. See H. J. Van Lonkhuijsen, 174

Hanai, T. See M. Ito, 325 Hänninen, O. See Y. Mälkki, 647

Hardmeier, R. M. See B. R. Hamaker, 461

Haridas Rao, P. See R. Sai Manohar, 619 Hartunian Sowa, S. M., and P. J. White. Characterization of starch isolated from oat groats with different amounts of lipid, 521

Hatcher, D. W. See B. A. Marchylo, 371

Hauber, R. J. See J. E. Fannon, 284 He, H., and R. C. Hoseney. Effect of the quantity of wheat flour protein on bread loaf volume, 17

.. Factors controlling gas retention in nonheated doughs, I R. R. Roach, and R. C. Hoseney. Effect of nonchaotropic salts on flour breadmaking properties, 366

Hill, A. S. See J. H. Skerritt, 110 Hisaki, K. See S. Aibara, 465

Hisaki, R. See A. Suzuki, 309
Hofer, P. J. See R. K. Newman, 240
Hojilla-Evangelista, M. P., L. A. Johnson, and D. J. Myers. Sequential extraction processing of flaked whole corn: Alternative corn fractionation technology for ethanol production, 643
Hoseney, R. C. See H. Dong, 132
\_\_\_\_\_. See H. He, 1, 17, 366
\_\_\_\_. See R. B. Malouf, 164, 169

See R. R. Roach, 574 See K. Shelke, 141, 145

baking. I. Effects of egg yolk and sugar ester, 89
Hruschka, W. R. See D. C. Slaughter, 428
Hsieh, H. M., Y. Pomeranz, and B. J. Swanson. Composition, cooking

time, and maturation of azuki (Vigna angularis) and common beans

(Phaseolus vulgaris), 244

Huang, M. L. See G. L. Rubenthaler, 334

Huckle, L. See K. Khan, 270, 686

Huebner, F. R., and C. S. Gaines. Relation between wheat kernel hardness, environment, and gliadin composition, 148 Hurburgh, C. R., Jr. See S. R. Fox, 191

Inagaki, T., and P. A. Seib. Firming of bread crumb with cross-linked

waxy barley starch substituted for wheat starch, 321

Inoue, Y., and W. Bushuk. Studies on frozen dough. II. Flour quality requirements for bread production from frozen dough, 423

Ito, M., S. Yoshikawa, K. Asami, and T. Hanai. Dielectric monitoring of gas production in fermenting bread dough, 325 Izydorczyk, M. S. See C. G. Biliaderis, 226

Jane, J.-L., and J.-F. Chen. Effect of amylose molecular size and amylopectin branch chain length on paste properties of starch, 60 , L. Shen, and F. Aguilar. Characterization of pejibaye starch,

L. Shen, J. Chen, S. Lim, T. Kasemsuwan, and W. K. Nip. Physical and chemical studies of taro starches and flours, 528

L. Wang, and C. C. Maningat. Preparation and properties

of small-particle corn starch, 280

A. Xu, M. Radosavljevic, and P. A. Seib. Location of amylose in normal starch granules. I. Susceptibility of amylose and amylopectin to cross-linking reagents, 405 Jayas, D. S. See N. D. G. White, 182

Johnson, L. A. See S. R. Fox, 191

See M. P. Hojilla-Evangelista, 643

See M. O. Racker, 66 See C. Wang, 43, 47 Jones, B. L. See K. Khan, 270

Kahlon, T. S., R. M. Saunders, R. N. Sayre, F. I. Chow, M. M. Chiu, and A. A. Betschart. Cholesterol-lowering effects of rice bran and rice bran oil fractions in hypercholesterolemic hamsters, 485

Kaneyama, M. See A. Suzuki, 309 Kasarda, D. D. See E. J.-L. Lew, 508 

Kassuba, A. See C. S. Gaines, 115, 120 Kavitha, R., and A. Chandrashekar. Content and composition of nonstarchy polysaccharides in endosperms of sorghums varying in hardness, 440

Kawakishi, S. See S. Nishibori, 160 Kawka, A., P. K. W. Ng, and W. Bushuk. Equivalence of high molecular weight glutenin subunits prepared by reversed-phase high-performance liquid chromatography and sodium dodecyl sulfate-polyacrylamide gel electrophoresis, 92

Kazemie, M., and W. Bushuk. NOTE: Presence of K proteins in developing wheat grain, 103 NOTE: Use of lithium chloride for the extraction of

and \_

flour proteins, 105 Khan, K., and L. Huckle. COMMUNICATION TO THE EDITOR: Use of multistacking gels in sodium dodecyl sulfate-polyacrylamide gel electrophoresis to reveal polydispersity, aggregation, and disaggregation of the glutenin protein fraction, 686

and B. L. Jones. Inheritance of gluten protein components of a high-protein hard red spring wheat line derived from Triticum turgidum var. dicoccoides—semipreparative RP-HPLC, gel electrophoresis, and amino acid composition studies, 270

See B. Aktan, 288 See J. Chen, 475, 481 See E. M. Magnus, 607 Khelifi, D. See G. Branlard, 677

Kim, C. S., and C. E. Walker. Effects of sugars and emulsifiers on starch gelatinization evaluated by differential scanning calorimetry, 212 and

Interactions between starches, sugars, and emulsifiers in high-ratio cake model systems, 206

Kim, S. See M. E. Carr, 70

Kim, S. S., and C. S. Setser. Wheat starch gelatinization in the presence of polydextrose or hydrolyzed barley β-glucans, 447

Kirkland, J. J. See B. A. Marchylo, 371 Kirleis, A. W. See D. L. Rom, 178

Klopfenstein, C. F. See R. K. Newman, 240 Knuckles, B. E., M. M. Chiu, and A. A. Betschart. β-Glucan-enriched fractions from laboratory-scale dry milling and sieving of barley and oats, 198

Knutson, C. A., and M. J. Grove. Variability of α-amylase synthesis in germinating maize, 436

in germinating matze, 430 Kohlstad, I. See B. R. Hamaker, 461 Kokini, J. L. See L. L. Chedid, 551 ——. See H. Madeka, 489 Kron, R. J., Sr. See E. T. Champagne, 152 Kruger, J. E. See B. A. Marchylo, 371 Kuzmicky, D. D. See E. J.-L. Lew, 508

Lai, C. S. See R. R. Roach, 574

Lambert, L. L. P., J. Gordon, and E. A. Davis. Water loss and structure development in model cake systems heated by microwave and convection methods, 303

Lang, C. E., E. K. Neises, and C. E. Walker. Effects of additives on flour-water dough mixograms, 587

Lanning, S. P., C. F. McGuire, and L. E. Talbert. NOTE: Parental blends as predictors of quality in spring wheat hybrids, 349

Lawton, J. W. Viscoelasticity of zein-starch doughs, 351

Lee, J. K. See M. H. Gomez, 275 Le Guernevé, C. See D. Bertrand, 257 Lehrfeld, J. See F. R. Dintzis, 577 Leisola, M. See A. Törrönen, 355

Lelievre, J. See H. Liu, 597

Lescano, M. See B. R. Hamaker, 461 Lew, E. J.-L., D. D. Kuzmicky, and D. D. Kasarda. Characterization of low molecular weight glutenin subunits by reversed-phase highperformance liquid chromatography, sodium dodecyl sulfate-polyacrylamide gel electrophoresis, and N-terminal amino acid sequencing,

Li, A., and C. E. Walker. NOTE: Dough temperature changes during mixing in a mixograph, 681

Liang, Y. T. See W. J. Lim, 233, 237
Lim, H., C. S. Setser, and J. V. Paukstelis. Comparisons of water mobility using <sup>17</sup>O nuclear magnetic resonance for four sugars: Glucose, maltose, maltotriose, and sucrose, 387

., and D. Sobczynska. 17O nuclear magnetic resonance studies on wheat starch-sucrose-water interactions with increasing temperature, 382

Lim, H. S., P. J. White, and K. J. Frey. Genotypic effects on β-glucan content of oat lines grown in two consecutive years, 262

Lim, S. See J.-L. Jane, 528

Lim, W. J., Y. T. Liang, and P. A. Seib. Cationic oat starch: Preparation and effect on paper strength, 237
\_\_\_\_\_\_\_, \_\_\_\_\_\_, and C. S. Rao. Isolation of oat starch from oat

flour, 233

Lin, W. D. A. See R. B. Malouf, 169 Lindahl, L., and A.-C. Eliasson. A comparison of some rheological properties of durum and wheat flour doughs, 30

.. Influence of added enzymes on the rheological properties of a wheat flour dough, 542

Litchfield, J. B. See R. Ruan, 13, 600

Liu, H., and J. Lelievre. Differential scanning calorimetric and rheological study of the gelatinization of starch granules embedded in a gel matrix,

Lookhart, G. See G. Branlard, 677
\_\_\_\_. See H. Dong, 132
Lorenz, K. See I. Al-Ruqaie, 472

See K. Delost-Lewis, 359 Lukow, O. M. See C. G. Biliaderis, 226

See K. R. Preston, 560

MacRitchie, F. See R. B. Gupta, 125 Madeka, H., and J. L. Kokini. Effect of addition of zein and gliadin on the rheological properties of amylopectin starch with low-tointermediate moisture, 489
Magnus, E. M., and K. Khan. Hydrophobic properties of reduced and

alkylated acetic acid-soluble glutenins-Fractionation by hydrophobic

interaction chromatography, 607 Malcolmson, L. J. See R. R. Matsuo, 27 Malkawi, H. I. See C. F. Morris, 467

Mälkki, Y., K. Autio, O. Hänninen, O. Myllymäki, K. Pelkonen, T. Suortti, and R. Törrönen. Oat bran concentrates: Physical properties of  $\beta$ -glucan and hypocholesterolemic effects in rats, 647 Malouf, R. B., and R. C. Hoseney. Wheat hardness. I. A method to

measure endosperm tensile strength using tablets made from wheat flour, 164

W. D. A. Lin, and R. C. Hoseney. Wheat hardness. II. Effect

of starch granule protein on endosperm tensile strength, 169
Mani, K., A.-C. Eliasson, L. Lindahl, and C. Trägärdh. Rheological
properties and breadmaking quality of wheat flour doughs made with

different dough mixers, 222

Maningat, C. C. See J.-L. Jane, 280

Marchylo, B. A., D. W. Hatcher, J. E. Kruger, and J. J. Kirkland.

Reversed-phase high-performance liquid column 271 wheat proteins using a new, highly stable column, 371 Marion, D. See D. Bertrand, 257

Marshall, W. E. Effect of degree of milling of brown rice and particle size of milled rice on starch gelatinization, 632

Martin, G. See P. Scheromm, 664

Matsumoto, H. See K. Hosomi, 89
Matsunoto, R. R., L. J. Malcolmson, N. M. Edwards, and J. E. Dexter.

A colorimetric method for estimating spaghetti cooking losses, 27 Mauromoustakos, A. See S. B. Andrews, 35 McCaig, T. N., J. G. McLeod, J. M. Clarke, and R. M. DePauw. NOTE: Measurement of durum pigment with a near-infrared instrument operating in the visible range, 671

McDonough, C. M. See M. H. Gomez, 275 McGuire, C. F. See S. P. Lanning, 349 McLeod, J. G. See T. N. McCaig, 671

Meuser, F. See A. Vollmar, 20 Mills, J. T. See N. D. G. White, 182 Mistry, A. H., and S. R. Eckhoff. Alkali debranning of corn to obtain corn bran, 202

Characteristics of alkali-extracted starch obtained from and \_ corn flour, 296

and \_ Dry milling and physical characteristics of alkali-

debranned yellow dent corn, 82

\_\_\_\_\_\_, M. P. Steinberg, and S. R. Eckhoff. Fractionation of high-lysine corn to produce edible by-products, 433

Moreira, R. G., and F. W. Bakker-Arkema. Grain dryer controls: A

review, 390

Morgan, B. See K. R. Preston, 560 Morris, C. F., G. A. Greenblatt, and H. I. Malkawi, COMMUNICATION TO THE EDITOR: Enhanced electrophoretic detection and isolation of friabilin, a starch granule protein, 467

Morrison, W. R. See R. F. Tester, 654 Myers, D. J. See M. P. Hojilla-Evangelista, 643

Myllymäki, O. See Y. Mälkki, 647

Natsuaki, O. See K. F. Finney, 347 Neises, E. K. See C. E. Lang, 587 Nelsen, T. C. See F. R. Dintzis, 577

Newman, C. W. See R. K. Newman, 240 Newman, R. K., C. F. Klopfenstein, C. W. Newman, N. Guritno, and P. J. Hofer. Comparison of the cholesterol-lowering properties of whole barley, oat bran, and wheat red dog in chicks and rats, 240

Ng, P. K. W. See L. Gao, 452 See A. Kawka, 92

Nip, W. K. See J.-L. Jane, 528

Nishibori, S., and S. Kawakishi. Effect of various sugars on the quality of baked cookies, 160

Nishio, K. See K. Hosomi, 89 Norris, K. H. See S. R. Delwiche, 107

See D. C. Slaughter, 428 Nussinovitch, A., M. S. Steffens, and P. Chinachoti. NOTE: Elastic properties of bread crumb, 678 See P. A. Rao, 613

O'Brien, L. See P. W. Gras, 254

Oliver, J. R., A. B. Blakeney, and H. M. Allen. Measurement of flour color in color space parameters, 546

Paszczynska, B. See Z. Czuchajowska, 516 Paukstelis, J. V. See H. Lim, 382, 387 Paulsen, M. R. See A. J. Peplinski, 397 Peleg, M. See S. Swyngedau, 217

Pelkonen, K. See Y. Mälkki, 647 Peplinski, A. J., M. R. Paulsen, and A. Bouzaher. Physical, chemical, and dry-milling properties of corn of varying density and breakage susceptibility, 397

Peterson, R. E., and W. J. Wolf. NOTE: Enhancement of highperformance liquid chromatography of soybean proteins by addition of sodium dodecyl sulfate, 101

Pitt, R. E. See S. R. Delwiche, 107 Pollak, L. See Y.-J. Wang, 328 Pomeranz, Y. See K. Addo, 6

See Z. Czuchajowska, 413, 516 See H. M. Hsieh, 244

See G. L. Rubenthaler, 334

See J. Szczodrak, 419, 626
Ponte, J. G., Jr. See A. Xu, 495, 502
Posner, E. S. See G. S. Ranhotra, 75
Preston, K. R., O. M. Lukow, and B. Morgan. Analysis of relationships between flour quality properties and protein fractions in a world wheat

collection, 560 Puhr, D. P., and B. L. D'Appolonia. Effect of baking absorption on bread yield, crumb moisture, and crumb water activity, 582

Radosavljevic, M. See J.-L. Jane, 405 Raeker, M. J., C. J. Bern, L. A. Johnson, and B. A. Glatz. Preservation of high-moisture maize by various propionate treatments, 66 Ranhotra, G. S., J. A. Gelroth, and B. K. Glaser. Lipidemic response

in rats fed flaxseed or sunflower oils, 623

and E. S. Posner. Total and soluble fiber content of air-classified white flour from hard and soft wheats, 75

Rao, C. S. See W. J. Lim, 233
Rao, P. A., A. Nussinovitch, and P. Chinachoti. Effects of selected surfactants on amylopectin recrystallization and on recoverability of bread crumb during storage, 613

Roach, R. R., C. S. Lai, and R. C. Hoseney. Effect of certain salts on bread loaf volume and on soluble nitrogen of wheat flour and nonfat dry milk slurries, 574

. See H. He. 366 Robert, P. See D. Bertrand, 257 Rodriguez, T. See B. R. Hamaker, 461

Rom, D. L., J. M. Shull, A. Chandrashekar, and A. W. Kirleis. Effects of cooking and treatment with sodium bisulfite on in vitro protein digestibility and microstructure of sorghum flour, 178 Rooney, L. W. See M. H. Gomez, 275

See S. O. Serna-Saldivar, 78 See N. E. Vivas, 673

Ruan, R., and J. B. Litchfield. Determination of water distribution and mobility inside maize kernels during steeping using magnetic resonance imaging, 13

and S. R. Eckhoff. Simultaneous and nondestructive measurement of transient moisture profiles and structural changes in corn kernels during steeping using microscopic nuclear magnetic resonance imaging, 600
Rubenthaler, G. L., Y. Pomeranz, and M. L. Huang. Steamed bread.

IV. Negative steamer-spring of strong flours, 334

Sai Manohar, R., and P. Haridas Rao. Use of a penetrometer for measuring rheological characteristics of biscuit dough, 619

Sanchez-Marroquin, A. See F. R. Del Valle, 156 Saunders, R. M. See T. S. Kahlon, 485

Sayre, R. N. See T. S. Kahlon, 485

Scheromm, P., G. Martin, A. Bergoin, and J.-C. Autran. Influence of nitrogen fertilization on the potential bread-baking quality of two wheat cultivars differing in their responses to increasing nitrogen supplies,

Schreuder, C. See H. J. Van Lonkhuijsen, 174

Schwartz, S. J. See S. E. Case, 401 Schwartzberg, H. G. See P. J. Wu, 567 Sears, R. G. See H. Dong, 132

Sehgal, S. See P. Gahlawat, 463 Seib, P. A. See T. Inagaki, 321 See J.-L. Jane, 405

See W. J. Lim, 233, 237

Sekhon, K. S., N. Singh, and R. P. Singh. Studies on the improvement of quality of wheat infected with Karnal bunt. I. Milling, rheological, and baking properties, 50

See N. Singh, 55 Serna-Saldivar, S. O., L. W. Rooney, and L. W. Greene. Effects of lime treatment on the bioavailability of calcium in diets of tortillas and beans: Bone and plasma composition in rats, 78

Setser, C. S. See A. M. Frye, 338

See S. S. Kim, 447 See H. Lim, 382, 387

Shelke, K., R. C. Hoseney, J. M. Faubion, and S. P. Curran. Age-related changes in the cake-baking quality of flour milled from freshly harvested soft wheat, 141

Age-related changes in the properties of and. batters made from flour milled from freshly harvested soft wheat, 145

Shelton, D. R. See J. Chen, 475, 481 Shen, L. See J.-L. Jane, 96, 280, 528 Shibanuma, K. See A. Suzuki, 309

Shogren, M. D. See H. Dong, 132 Shull, J. M. See D. L. Rom, 178

Siebenmorgen, T. J. See S. B. Andrews, 35 Sinda, E. See K. Autio, 409 Singh, N., K. S. Sekhon, A. K. Srivastava, and P. P. Gupta. Studies on the improvement of quality of wheat infected with Karnal bunt. II. Nutritional and biological effects, 55

See K. S. Sekhon, 50

Singh, R. P. See K. S. Sekhon, 50 Skerritt, J. H., and A. S. Hill. NOTE: How "free" is "gluten free"? Relationship between Kjeldahl nitrogen values and gluten protein content for wheat starches, 110

Slaughter, D. C., K. H. Norris, and W. R. Hruschka. Quality and classification of hard red wheat, 428

Sobczynska, D. See H. Lim, 382 Srivastava, A. K. See N. Singh, 55 Stanley, K. D. See M. E. Carr, 70

Steffens, M. S. See A. Nussinovitch, 678 Steinberg, M. P. See A. H. Mistry, 433 Stringfellow, A. C. See Y. V. Wu, 188

Strong III, F. C., and A. M. A. Duarte. A room-temperature, rapid method for the determination of protein in wheat and other grains by the biuret reaction, 659

Suortti, T. See Y. Mälkki, 647

Suzuki, A., M. Kaneyama, K. Shibanuma, Y. Takeda, J. Abe, and S. Hizukuri. Characterization of lotus starch, 309

Swanson, B. G. See H. M. Hsieh, 244

Swyngedau, S., and M. Peleg. Characterization and prediction of the compressive stress-strain relationship of layered arrays of spongy baked goods, 217

Symons, S. J., and J. E. Dexter. Estimation of milling efficiency: Prediction of flour refinement by the measurement of pericarp fluorescence, 137 Szczodrak, J., Z. Czuchajowska, and Y. Pomeranz. Characterization and estimation of barley polysaccharides by near-infrared spectroscopy. II.

Estimation of total β-D-glucans, 419
and Y. Pomeranz. Starch-lipid interactions and formation of resistant starch in high-amylose barley, 626

See Z. Czuchajowska, 413

Takeda, Y. See A. Suzuki, 309

Talbert, L. E. See S. P. Lanning, 349 Tester, R. F., and W. R. Morrison. Swelling and gelatinization of cereal starches. III. Some properties of waxy and normal nonwaxy barley starches, 654

Tkachuk, R. See G. S. Chauhan, 85

Törrönen, A., M. Leisola, and S. Haarasilta. Inhibition of rye α-amylase activity by barley α-amylase inhibitor, 355 Törrönen, R. See Y. Mälkki, 647

Trägärdh, C. See K. Mani, 222

Tribelhorn, R. See K. Delost-Lewis, 359

Umbach, S. L., E. A. Davis, J. Gordon, and P. T. Callaghan. Water self-diffusion coefficients and dielectric properties determined for starchgluten-water mixtures heated by microwave and by conventional methods, 637

Valdivia, R. See B. R. Hamaker, 461 Valles, C. See B. R. Hamaker, 461

Van Lonkhuijsen, H. J., R. J. Hamer, and C. Schreuder. Influence of specific gliadins on the breadmaking quality of wheat, 174 Vivas, N. E., R. D. Waniska, and L. W. Rooney. NOTE: Effects on

proteins in sorghum, maize, and pearl millet when processed into acidic and basic Tô, 673

Vollmar, A., and F. Meuser. Influence of starter cultures consisting of lactic acid bacteria and yeasts on the performance of a continuous sourdough fermenter, 20

Walker, C. E. See C. S. Kim, 206, 212

See C. E. Lang, 587 See A. Li, 681

See A. R. Wooding, 249

See Y. Yin, 592

Wang, C., and L. A. Johnson. Wet-milling characteristics of propionatetreated high-moisture maize. I. Yields and compositions of products, 43

.. Wet-milling characteristics of propionate-treated highmoisture maize. II. Qualities of starch and gluten, 47

Wang, L. See J.-L. Jane, 280

Wang, Y.-J., P. White, and L. Pollak. Thermal and gelling properties of maize mutants from the OH43 inbred line, 328

Waniska, R. D. See M. H. Gomez, 275 See N. E. Vivas, 673

Watanabe, K. See S. Aibara, 465

Werner, W. E., A. E. Adalsteins, and D. D. Kasarda. Composition of high-molecular-weight glutenin subunit dimers formed by partial reduction of residue glutenin, 535

White, N. D. G., D. S. Jayas, J. T. Mills, and B. L. Dronzek. Effects of canola oil or white mineral oil at dust suppressant levels on the storage characteristics of wheat, 182

White, P. J. See S. M. Hartunian Sowa, 521

mechanisms (mobile vs. fixed-bowl) for the 35- and 10-gram mixographs, 249

Wu, P. J., and H. G. Schwartzberg. Popping behavior and zein coating

of popcorn, 567
Wu, Y. V. Corn hardness as related to yield and particle size of fractions from a micro hammer-cutter mill, 343

and A. C. Stringfellow. Air classification of flours from wheats with varying hardness: Protein shifts, 188

Xu, A., O. K. Chung, and J. G. Ponte, Jr. Bread crumb amylograph studies. I. Effects of storage time, shortening, flour lipids, and surfactants, 495

\_\_\_\_, J. G. Ponte, Jr., and O. K. Chung. Bread crumb amylograph studies. II. Cause of unique properties, 502

See J.-L. Jane, 405

Yin, Y., and C. E. Walker. Pentosans from gluten-washing wastewater: Isolation, characterizations, and role in baking, 592

Yoon, K. J. See M. E. Barr, 70 Yoshikawa, S. See M. Ito, 325

## Subject Index

Acknowledgment of reviewers, vi

Additives, effect on flour-water dough mixographs (Lang et al), 587

Air classification, of wheat flour, relation of hardness to protein content, yield, and protein shift (Wu and Stringfellow), 188

debranning, to obtain corn bran (Mistry and Eckhoff), 202

extracted starch, from corn flour, characteristics (Mistry and Eckhoff),

Alkylresorcinols, in extruded cereal brans (Al-Ruqaie and Lorenz), 472 Alveograph, characteristics; effects of lipids, emulsifiers, defatting of flours (Addo and Pomeranz), 6

Amaranth, in infant formula, nitrogen balance (Del Valle et al), 156 Amino acid, composition

-of carbohydrate-containing protein (Chen et al), 475

of gliadin fractions (Khan et al), 270

Amino acid sequencing, sequences of and cysteines in LMW-glutenin subunits (Lew et al), 508

inhibition of rye and barley (Törrönen et al), 355

-variability of synthesis in germinating maize (Knutson and Grove),

Amylopectin, recrystallization, surfactant effect (Rao et al), 613 Amylose

colorimetric determination (Tester and Morrison), 654

interaction with lipid, cause of unique properties of bread crumb amylogram (Ku et al), 502

location in starch granules (Jane et al), 405

### Baking

absorption, effect on bread yield, crumb moisture, and crumb water activity (Puhr and D'Appolonia), 582

bread: effect of pentosans from gluten-washing water (Yin and Walker), 592; rheological properties and breadmaking quality of doughs made with different mixers (Mani et al), 222

-crumb of bread characterized by video image analysis (Bertrand et al), 257

effects of egg yolk and sugar ester on frozen dough (Hosomi et al),

gas retention in nonheated doughs (He and Hoseney), 1

instrumental measurement of cookie hardness: application to product quality variables (Gaines et al), 120; assessment of methods (Gaines et al), 115

potassium bromate analysis in flour, dough, and bread (De Stefanis), 683

-protein quantity effect on bread loaf volume (He and Hoseney), 17 quality: flours milled from freshly harvested soft wheats (Shelke et al), 141; relationship to protein fractions (Preston et al), 560

sensory characteristics and optimization of sugar-substituted cakes (Frye and Setser), 338

use of electrical resistance oven for freshly harvested soft wheat batters (Shelke et al), 145

white pan and whole wheat breads, comparison of hard red and white wheats (Chang and Chambers), 556

#### Barley

α-amylase activity inhibition (Törrönen et al), 355

cholesterol-lowering properties (Newman et al), 240

—dry milling and sieving for β-glucan enrichment (Knuckles et al), 198

β-glucan content and viscosities of (Bhatty), 469

β-D-glucans in (Szczodrak et al), 419

high-amylose, starch-lipid interactions (Szczodrak and Pomeranz), 626 hydrolyzed β-glucan effects on wheat starch gelatinization (Kim and Setser), 447

-near-infrared reflectance spectra (Czuchajowska et al), 413

-starch, waxy-nonwaxy, granule dimensions and swelling factors (Tester and Morrison), 654

Beans, azuki and common, composition, cooking time, and maturation of (Hsieh et al), 244

Bioavailability, calcium, in tortillas (Serna-Saldivar et al), 78

Biuret reaction, for rapid determination of protein in grains (Strong and Duarte), 659

Bran, phytate content as related to kernel size, environmental influences, and soft spring cultivars (Dintzis et al), 577

Bread and breadmaking

-baking absorption effect on bread crumb moisture, yield, and crumb water activity (Puhr and D'Appolonia), 582

-bread crumb amylogram, cause of unique properties (Ku et al), 502

-bread crumb: characterized by video image analysis (Bertrand et al),

257; during storage, surfactant effect (Rao et al), 613; elastic properties of (Nussinovitch et al), 678

bread crust and crumb, aroma and flavor (Chang and Chambers), 556 -compressive stress-strain of crumbs, characterization and prediction

(Swyngedau and Peleg), 217

copper (II) vs. zinc inorganic salts as oxidizers in (Finney et al), 347 crumb amylograph studies, effects of storage time, shortening, flour lipids, and surfactants (Ku et al), 495

effects of wheat proteins (Scheromm et al), 664

firming of crumb containing waxy barley starch (Inagaki and Seib), 321

influence of specific gliadins (Van Lonkhuijsen et al), 174

monitoring of gas production in bread dough (Ito et al), 325

-nonchaotropic salt effect on (He et al), 366

potassium bromate analysis in (De Stefanis), 683

properties, effects of high-voltage electric field treatment (Aibara et al), 465

protein composition and breadmaking quality in wheat (Dong et al),

-psyllium effect on (Czuchajowska et al), 516

quality and rheological properties of, using different dough mixers (Mani et al), 222

-reduction of phytic acid in whole meal (Fretzdorff and Brümmer), 266

salt effect on interaction of milk proteins and wheat flour proteins (Roach et al), 574

-steamed, oven spring of (Rubenthaler et al), 334

Brown rice, stabilizing to lipolytic hydrolysis by ethanol vapors (Champagne and Hron), 152

age-related changes in batter properties (Shelke et al), 145

baking quality of, age-related changes in, freshly harvested soft wheats (Shelke et al), 141

compressive stress-strain of crumbs, characterization and prediction (Swyngedau and Peleg), 217

model systems, water loss and structure development in, microwave and convection heating (Lambert et al), 303

psyllium effect on (Czuchajowska et al), 516

reduced-calorie, layer-sensory and optimization (Frye and Setser), 338 starch, sugar, and emulsifier interaction (Kim and Walker), 206

Calcium, bioavailability in tortillas (Serna-Saldivar et al), 78

Carbohydrate

in glycoprotein of gluten (Chen et al), 475 structural characterization of glycopeptide of wheat gluten (Chen et al), 481

Chlorination, effect on flours milled from freshly harvested soft wheats (Shelke et al), 141

Cholesterol

-effect: of barley, oat bran, and wheat in chicks and rats (Newman et al), 240; of rice bran, defatted rice bran plus rice bran oil, and rice bran oil fractions-wax, gum, degummed-dewaxed oil (Kahlon et al), 485

reduction by oat bran and concentrates (Mälkki et al), 647

Chromatography

hydrophobic interaction on phenyl-Sepharose CL-4B (Magnus and Khan), 607

-semipreparative RP-HPLC of gliadin (Khan et al), 270

-size-exclusion: of gluten proteins (Chen et al), 475; and ion-exchange of glycopeptide from gluten (Chen et al), 481

effect of various sugars on quality (Nishibori and Kawakishi), 160

instrumental measurement of cookie hardness: application to product quality variables (Gaines et al), 120; assessment of methods (Gaines et al), 115

Corn (see also Maize)

alkali-debranned yellow dent, dry milling and physical characteristics of (Mistry and Eckhoff), 82

bran, alkali debranning to obtain (Mistry and Eckhoff), 202

breakage susceptibility, density, stress cracked kernels, hardness, drymilling response, and proximate analysis (Peplinski et al), 397

-calcium bioavailability in tortillas (Serna-Saldivar et al), 78

-hardness, related to particle size and yield of fractions from a micro hammer-cutter mill (Win), 343

-high-lysine, fractionation to produce edible by-products (Mistry et al),

-kernels, determination of water distribution and mobility in, using MRI (Ruan and Litchfield), 13

- -moisture measurement using MRI (Ruan et al), 600
- —sequential extraction processing using ethanol (Hojilla-Evangelista et al), 643
- -starch changes during tortilla processing (Gomez et al), 275
- -sweet, zeins in (Wilson), 113

Dietary fiber, nonstarch polysaccharide and lignin values compared (Flint and Camire), 444

Dielectric properties, of starch-gluten-water system (Umbach et al), 637 Differential scanning calorimetry

- —comparisons for glucose, maltose, maltotriose, and sucrose on starch onset temperature (Lim et al), 387
- of flour with repeated heating and cooling (Ku et al), 502
   of gelatinization of starch granules (Liu and Lelievre), 597
- of high-amylose barley, starch-lipid interaction (Szczodrak and Pomeranz), 626
- -for sugar and emulsifier effect on starch gelatinization (Kim and Walker), 212
- Dough
- breadmaking properties, effects of high-voltage electric field treatment (Aibara et al), 465
- -dielectric monitoring of dough expansion (Ito et al), 325
- —frozen: effects of egg yolk and sugar ester (Hosomi et al), 89; rheology and yeast viability (Autio and Sinda), 409
- —frozen bread, storage and freeze-thaw cycle effect and flour quality on baking and rheological properties (Inoue and Bushuk), 423
- mixing properties: effect of protein and lipoprotein fractions (Bekes and Gras), 229; heritability in early generations (Gras and O'Brien), 254
- -potassium bromate analysis in (De Stefanis), 683
- --reduction of phytic acid in model dough systems as affected by pH and temperature (Fretzdorff and Brümmer), 266
- --rheological characteristics of biscuit dough, effect of mixing time and amount of water (Sai Manohar and Haridas Rao), 619
- —rheological properties and breadmaking quality with different dough mixers (Mani et al), 222
- -temperature, during mixing (Li and Walker), 681
- -treated with low concentrations of reducing agent (Khan and Huckle), 686
- viscoelastic properties of wheat flour doughs with added enzymes and at raised temperatures (Lindahl and Eliasson), 542
- -viscoelasticity of zein-starch mixtures (Lawton), 351
- Dry milling, of alkali-debranned yellow dent corn (Mistry and Eckhoff),
- Drying, effect on quality and biochemical components of pasta (Aktan and Khan), 288

# Elastic properties, of bread crumb (Nussinovitch et al), 678 Electrophoresis

- -acid and SDS separation of wheat proteins (Branlard et al), 677
- of fractions of glutenin from hydrophobic interaction chromatography (Magnus and Khan), 607
- —of glutenin in relation to structure, SDS-PAGE modified procedure (Khan and Huckle), 686
- -of high-temperature dried pasta (Aktan and Khan), 288
- —one- and two-dimensional SDS-PAGE of partially reduced glutenin (Werner et al), 535
- -of RP-HPLC gliadin fractions (Van Lonkhuijsen et al), 174
- Emulsitiers
  —effect: on alveograph characteristics (Addo and Pomeranz), 6; on bread crumb amylogram and crumb firmness (Ku et al), 495; on starch gelatinization (Kim and Walker), 212
- -interaction with starches and sugars in high-ratio cake model systems (Kim and Walker), 206
- Ethanol, recovery from corn during sequential extraction processing (Hojilla-Evangelista et al), 643
- Environment
- —effect on bran phytate content and flour quality and milling parameters (Dintzis et al), 577
- —and nitrogen fertilization, effect on baking quality (Scheromm et al), 664
- Enzymes, addition of  $\alpha$ -amylases and mixtures of  $\alpha$ -amylases and proteases to wheat flour doughs (Lindahl and Eliasson), 542
- Ethanol, vapors, for lipolytic hydrolysis to stabilize brown rice (Champagne and Hron), 152
- Extrusion, of corn meal, corn starch, wheat flour, and wheat starch, gelatinization and textural properties (Case et al), 401

#### Fiber

-soluble, from oat bran and concentrate, physical properties and hypo-

- cholesterolemic effects in rats (Mälkki et al), 647
- -total and soluble, in air-classified white flour (Ranhotra et al), 75 Flour
- -air-classified white, fiber content of (Ranhotra et al), 75
- -color, measurement in color space parameters (Oliver et al), 546
- -corn, alkali-extracted starch obtained from (Mistry and Eckhoff), 296 -defatted, alveograph characteristics of (Addo and Pomeranz), 6
- -nonchaotropic salt effect in breadmaking (He et al), 366
- -oat, isolation of oat starch from (Lim et al), 233
- —potassium bromate analysis in (De Stefanis), 683
- procedure to make tablets of to measure endosperm tensile strength (Malouf and Hoseney), 164
- -refinement prediction; measurement of pericarp fluorescence (Symons and Dexter), 137
- -taro, physicochemical studies (Jane et al), 528
- -wheat, functional properties and protein composition (Gupta et al), 125

#### Fractionation

- -of barley and oats by dry milling and sieving, yield and  $\beta$ -glucan and starch content of fractions (Knuckles et al), 198
- of carbohydrate-containing proteins of gluten (Chen et al), 475
- —of glutenin by hydrophobic interaction chromatography (Magnus and Khan), 607
- -of glycoprotein from gluten (Chen et al), 481

# Gas retention, in nonheated doughs (He and Hoseney), 1 Gelatinization

- of extruded starch-based materials, relationship to textural properties (Case et al), 401
- —properties of starches from maize mutants (Wang et al), 328
- Gliadin
  —effect on rheological properties of amylopectin starch (Madeka and
- Kokini), 489
  —in relation to: baking quality (Van Lonkhuijsen et al), 174; kernel
- hardness, environment, and spike position (Huebner and Gaines), 148  $\beta$ -Glucan
- —content in barley and oat fractions produced by dry milling and sieving (Knuckles et al), 198
- -measurement in oat lines grown two years (Lim et al), 262
- —from oat bran and concentrates, physical properties and hypocholesterolemic effects in rats (Mälkki et al), 647
- —digestion with pronase enzyme (Chen et al), 481
- —free, Kjeldahl nitrogen values and gluten protein content for wheat starches (Skerritt and Hill), 110
- -metal-complexing agent effect on water binding by (Clements), 315
- -characterization of LMW-glutenin subunits (Lew et al), 508
- —dimers of HMW-glutenin subunits formed by partial reduction (Werner et al), 535
- —equivalence of subunits prepared by RP-HPLC and SDS-PAGE (Kawka et al), 92
- -fractionation by hydrophobic interaction chromatography (Magnus and Khan), 607
- -HMW and LMW (Branlard et al), 677
- -SDS-PAGE of unreduced protein (Khan and Huckle), 686
- -structure polymeric model in relation to rheological properties (Gao et al), 452

#### Grain

- antinutrients as affected by oven heating (Gahlawat and Sehgal), 463
   dryers, in-bin and continuous-flow (Moreira and Bakker-Arkema), 390
- Guideline for the preparation and review of papers reporting sensory evaluation data, v

#### Hardnes

- —of cookies: instrumental measurement of, application to product quality variables (Gaines et al), 120; instrumental measurement of, assessment of methods (Gaines et al), 115
- -of corn, related to yield and particle size of fractions (Wu), 343
- —of wheat: measuring endosperm tensile strength of tablets made from wheat flour (Malouf and Hoseney), 164; relating to gliadin composition (Huebner and Gaines), 148; starch granule protein effect on (Manouf et al), 169

#### HPLC

- -characterization by lotus starch by (Suzuki et al), 309
- -enhancement by addition of SDS (Peterson and Wolf), 101

#### IEF, zeins in sweet corn (Wilson), 113

#### Image analysi

-bread crumb, characterization of texture (Bertrand et al), 257

-to determine starch particle sizes (Jane et al), 280 Instructions to authors, iii

Lactic acid, bacteria, in starter cultures, effect on sourdough (Vollmar and Meuser), 20

Lignin, recovery during nonstarch polysaccharide analysis (Flint and Camire), 444

Lipase, effect on lipid stability in oats (Ekstrand et al), 379

Lipidic response, in rats fed flaxseed or sunflower oils (Ranhotra et al), 623

#### Lipids

-deterioration related to lipase activity in oats (Ekstrand et al), 379

-effect on alveograph characteristics (Addo and Pomeranz), 6

-flour, effect on bread crumb amylogram and crumb firmness (Ku et al), 495

 interaction with starch, cause of unique properties of bread crumb amylogram (Ku et al), 502

#### Maize (see also Corn)

- —high-moisture, preservation by propionate treatments (Raeker et al), 66
- -thermal and gelling properties of starches from mutants (Wang et al), 328
- —variability of  $\alpha$ -amylase synthesis in germinating (Knutson and Grove), 436
- wet-milling characteristics of: product yield and composition (Wang and Johnson), 43; proximate composition and physical property relation (Fox et al), 191; starch and gluten qualities (Wang and Johnson), 47 Methods

-estimating pasta cooking loss (Matsuo et al), 27

- —instrumental measurement of cookie hardness: application to product quality variables (Gaines et al), 120; assessment of methods (Gaines et al), 115
- -to monitor dough expansion dielectrically (Ito et al), 325
- room-temperature, rapid, for protein determination in grains (Strong and Duarte), 659

#### Microwave heating

- -model cake systems (Lambert et al), 303
- -starch-gluten-water system (Umbach et al), 637

Millet, puffing quality (Delost-Lewis et al), 359

#### Milling

- --to achieve desirable rheological behavior (Lindahl and Eliasson), 30 --of brown rice, thermal properties of starch in milled fractions (Marshall),
- -characteristics of Canadian wheats (Symons and Dexter), 137
- -degree evaluated by McGill rice miller (Andrews et al), 35
- —sequential extraction processing of corn (Hojilla-Evangelista et al), 643 Mixograph
- application to early generation selection for dough strength (Gras and O'Brien), 254
- -comparison of 35 gram vs. 10 gram (Wooding and Walker), 249
- -flour-water dough, additive effect on (Lang et al), 587
- -protein relationship with (Dong et al), 132
- temperature changes during mixing (Li and Walker), 681
- Moisture, in starch and microcrystalline cellulose as measured using NIR spectroscopy (Delwiche et al), 107

NIR, sensitivity to moisture content and water activity (Delwiche et al), 107

### NIR spectroscopy

- -of barley β-D-glucans (Szczodrak et al), 419
- -of barleys, starches, and β-D-glucans (Czuchajowska et al), 413

#### Nitrogen

- -balance, in infant formula (Del Valle et al), 156
- -digestibility of puffed millet (Delost-Lewis et al), 359

#### NMR

- —comparisons of water mobility for four sugars in wheat starch-water system (Lim et al), 387
- -pulsed gradient spin-echo, starch-gluten-water system (Umbach et al), 637
- -wheat starch-sucrose-water interactions (Lim et al), 382
- NMR imaging, water distribution and mobility inside corn kernels during steeping (Ruan and Litchfield), 13

#### Oat bran

- -cholesterol-lowering properties (Newman et al), 240
- —concentrates of, physical properties of  $\beta$ -glucan and hypocholesterolemic effects in rats (Mälkki et al), 647

#### Oats

-characterization of starch from groats with different amounts of lipid

- (Hartunian Sowa and White), 521
- —dry milling and sieving for  $\beta$ -glucan enrichment (Knuckles et al), 198
- -β-glucan content in different genotypes (Lim et al), 262
- -lipase activity (Ekstrand et al), 379

#### Oil

- —canola oil, mineral oil, dust suppression, storage, HRS wheat (White et al), 182
- -fatty acid profile of Inca peanut (Hamaker et al), 461
- -flaxseed or sunflower, lipidemic response in rats (Ranhotra et al), 623

Particle size, determination of small-particle corn starch (Jane et al), 280 Pasta

- -estimating cooking loss colorimetrically (Matsuo et al), 27
- -high-temperature effect on dried (Aktan and Khan), 288

#### Pentosans

- -in flours of 1B/1R translocation wheats (Biliaderis et al), 226
- -water-soluble, from gluten-washing water (Yin and Walker), 592
- Phytate, bran content, relationships to milling and flour quality parameters (Dintzis et al), 577
- Phytic acid, reduction during breadmaking of whole-meal breads (Fretzdorff and Brümmer), 266
- Polydextrose, effects of fractions on wheat starch gelatinization (Kim and Setser), 447
- Popcorn, popping behavior and zein coating of (Wu and Schwartzberg), 567

Porridge, stiff, acidic and basic (Vivas et al), 673

Propionic acid, for preservation of high-moisture maize (Raeker et al), 66

#### Protein

- -content and strength effect on oven spring of steamed bread (Rubenthaler et al), 334
- -fractionation by RP-HPLC and Osborne procedure (Khan et al), 270
- -friabilin starch granule (Morris et al), 467
- -glutenin oligomers and imers of subunits (Gao et al), 452
- -in grains, rapid determination (Strong and Duarte), 659
- influence on rheological properties of starches (Chedid and Kokini),
   551
- -K protein in developing wheat grain (Kazemie and Bushuk), 103
- -quality of Inca peanut (Hamaker et al), 461
- -quantity effect on bread loaf volume (He and Hoseney), 17
- -quinoa seed fractions (Chauhan et al), 85
- relationships between composition and functional properties of wheat flours (Gupta et al), 125
- -sorghum flour, in vitro digestibility of after cooking and treatment with sodium bisulfite (Rom et al), 178
- --in sorghum, maize, and pearl millet for stiff porridge (Vivas et al), 673
- -structure of disulfide-linked HMW-glutenin subunits (Werner et al), 535
- —of wheat flour, solubility in LiCl solution (Kazemie and Bushuk), 105 Psyllium, effects on bread staling and dough mixing (Czuchajowska et al), 516

Quinoa, nutrients and antinutrients (Chauhan et al), 85

#### Rheological properties

- —doughs, relation to breadmaking quality and mixing equipment (Mani et al), 222
- —effect of mixing time and amount of water on biscuit dough (Sai Manohar and Haridas Rao), 619

### Rheology

- -amylose, amylopectin plus protein systems (Chedid and Kokini), 551
- -frozen storage effect on (Autio and Sinda), 409
- -of gelatinization of starch granules (Liu and Lelievre), 597
- oscillatory measurements on wheat flour doughs (Lindahl and Eliasson),
   542
- -viscoelastic properties of durum and wheat flour doughs (Lindahl and Eliasson), 30

#### Rice

- -head, yield evaluated by McGill rice miller (Andrews et al), 35
- -starch thermal properties in brown and milled (Marshall), 632
- Rice bran, cholesterol-lowering in hamsters (Kahlon et al), 485 RP-HPLC
- -equivalence of HMW subunits prepared by (Kawka et al), 92
- —of gliadins: of high-temperature dried pasta (Aktan and Khan), 288; relating to kernel hardness (Huebner and Gaines), 148; in relation to baking quality (Van Lonkhuijsen et al), 174
- -separation of reduced and pyridylethylated fractions of LMW-glutenin subunits (Lew et al), 508
- -of wheat proteins using highly stable column (Marchylo et al), 371

Rye, α-amylase activity inhibition (Törrönen et al), 355

Rye bran, extruded, alkyresorcinols in (Al-Ruqaie and Lorenz), 472

Salt, effect on interaction of milk proteins and wheat flour proteins (Roach

Scanning electron microscopy

of sorghum flour, effects of cooking and treatment with sodium bisulfite (Rom et al), 178

of starch granules (Fannon et al), 284

- of starch paste and granule remnants (Fannon and BeMiller), 456 SDS-PAGE
- -analysis of unreduced glutenin by modified stacking gel procedure (Khan and Huckle), 686

-enhanced detection and isolation of friabilin (Morris et al), 467

equivalence of HMV subunits prepared by (Kawka et al), 92 of fractions of glutenin from hydrophobic interaction chromatography (Magnus and Khan), 607

-of high-temperature dried pasta (Aktan and Khan), 288

one-dimensional analysis of reduced and pyridylethylated fractions of LMW-glutenin subunits (Lew et al), 508

RP-HPLC of soybean proteins (Peterson and Wolf), 101

-two-dimensional analysis (unreduced vs. fully reduced proteins) of partially reduced glutenin (Werner et al), 535

-zeins in sweet corn (Wilson), 113

Sorghum

- -endosperm flour, nonstarchy polysaccharides and hardness (Kavitha and Chandrashekar), 440
- -flour, effects of cooking and treatment with sodium bisulfite on digestibility and microstructure (Rom et al), 178

Sourdough, fermenter, performance influenced by starter cultures (Vollmar and Meuser), 20

Soybean, proteins; enhanced HPLC with SDS (Peterson and Wolf), 101 Starch

- amylopectin, zein and gliadin effect on rheological properties of (Madeka and Kokini), 489
- amylose and amylopectin structures affecting paste properties (Jane and Chen), 60
- -barley, waxy, nonwaxy, granule dimensions and swelling factors (Tester and Morrison), 654

changes during tortilla processing (Gomez et al), 275

- -characterization from groats with different amounts of lipid (Kartunian Sowa and White), 521
- -corn, paste and granule remnants, scanning electron microscopy of (Fannon and BeMiller), 456

detection of water within using NIP spectroscopy (Delwiche et al),

-gelatinization: in brown and milled rice and pulverized rice kernels (Marshall), 632; DSC and rheology for (Liu and Lelievre), 597; in presence of bulking agents (Kim and Setser), 447; sugar and emulsifier effect on (Kim and Walker), 212

gelatinization in presence of bulking agents (Kim and Setser), 447

graft polymerization by reactive extrusion (Carr et al), 70

granular structure of (Jane et al), 405

granule protein; effect on wheat hardness (Manouf et al), 169

interaction: with sucrose and water studied by NMR (Lim et al), 382; with sugars by DSC and NMR studies (Lim et al), 387; with sugars and emulsifiers in high-ratio cake model systems (Kim and Walker), 206

lotus, characterization of (Suzuki et al), 309

near-infrared reflectance spectra (Czuchajowska et al), 413

oat: cationic, preparation and effect on paper strength (Lim et al), 237; isolation from oat flour (Lim et al), 233

-pejibaye, characterization (Jane et al), 96

- -rheological behavior of wheat flour doughs caused by damaged starch (Lindahl and Eliasson), 30
- -rheological properties of amylose, amylopectin plus protein systems (Chedid and Kokini), 551
- -small-particle corn, preparation and properties (Jane et al), 280

-surface pores of granules (Fannon et al), 284

-taro, physicochemical studies (Jane et al), 528

- -thermal and gelling properties of maize mutants (Wang et al), 328 -wheat, relation between Kjeldahl nitrogen values and gluten protein content (Skerritt and Hill), 110
- Steeping, of corn kernels, MRI for moisture measurement (Ruan et al),

Storage, effect of canola oil or mineral oil on wheat storage (White et al), 182

-effect: on cookie quality (Nishibori and Kawakishi), 160; on starch and water interactions studied by NMR (Lim et al), 382; on starch gelatinization (Kim and Walker), 212

influence on water mobility (Lim et al), 387

-interaction with starches and emulsifiers in high-ratio cake model systems (Kim and Walker), 206

Surfactants, effect

on anylopectin recrystallization and bread crumb recoverability (Rao et al), 613

on frozen dough (Hosomi et al), 89

Tannins, quinoa seed fractions (Chauhan et al), 85

Texture, of extruded starch-based materials, relationship to extent of gelatinization (Case et al), 401

Tortillas

-calcium bioavailability in (Serna-Saldivar et al), 78

-corn, effect of processing on starch (Gomez et al), 275

Triticale bran, extruded, alkyresorcinols in (Al-Ruqaie and Lorenz), 472

Viscosity, batter, age-related changes, in freshly harvested soft wheats (Shelke et al), 145

Wet milling

- of maize, grain proximate composition and physical property relations (Fox et al), 191
- of propionate-treated high-moisture maize: product yield and composition (Wang and Johnson), 43; starch and gluten qualities (Wang and Johnson), 47

Wheat

- -1B/1R translocation, pentosans in flours of (Biliaderis et al), 226
- -cultivar and environmental effect on quality (Scheromm et al), 664 durum, measurement of pigment by VIS/NIR (McCaig et al), 671
- durum and wheat flour doughs compared (Lindahl and Eliasson), 30 effect of oil application for dust suppression on storage characteristics,

baking and milling, and insecticides in HRS wheat (White et al), 182 estimating pasta cooking loss (Matsuo et al), 27

-flour, functional properties and protein composition (Gupta et al), 125 -flour protein solubility in LiCl solution (Kazemie and Bushuk), 105 -friabilin starch granule protein (Morris et al), 467

-hard red, classification and quality of (Slaughter et al), 428

-hard red and hard white winter, compared in bread (Chang and Chambers), 556

-hard and soft, air-classified white flour from, fiber content of (Ranhotra et al), 75

-hardness: measured by protein content, yield, and protein shift after air classification (Wu and Stringfellow), 188; measuring endosperm tensile strength of tablets made from wheat flour (Malouf and Hoseney), 164; starch granule protein effect on (Manouf et al), 169

hybrids, parental blends as predictors of quality (Lanning et al), 349

K proteins in developing grain (Kazemie and Bushuk), 103

Karnal bunt infected: milling, rheological, and baking properties (Sekhon et al), 50; nutritional and biological effects (Singh et al), 55 protein: glutenins, HMV and LMW, D-zone proteins (Branlard et al), 677; of high protein line (Khan et al), 270; RP-HPLC for analysis

with highly stable column (Marchylo et al), 371 -protein fractions, relationship to quality (Preston et al), 560

-relation of gliadins to kernel hardness (Huebner and Gaines), 148 -selection for mixing properties, use of 2-gram mixograph for (Gras and O'Brien), 254

-storage proteins, glutenin structure (Gao et al), 452

-structure and arrangement of glutenin protein subunits (Werner et al), 535

Wheat bran, extruded, alkyresorcinols in (Al-Ruqaie and Lorenz), 472

- -activity, frozen storage effect in dough (Autio and Sinda), 409
- -in starter cultures, effect on sourdough (Vollmar and Meuser), 20

- -coating, of popcorn (Wu and Schwartzberg), 567
- effect on rheological properties of amylopectin starch (Madeka and

in sweet corn (Wilson), 113

-viscoelasticity of composite flour doughs containing starch and dibutyl tartrate (Lawton), 351

